



THE ORTHOPEDIC PARTNERS  
AN RCM CLINIC

PARK CITY • HEBER CITY • SALT LAKE CITY

M. Daniel Hatch, M.D.

[www.utahshoulder.com](http://www.utahshoulder.com)

## Labral Repair Protocol

### Phase 1

Sling

- 1) Full time use for first 6 weeks
- 2) Discontinue after week 6

Weeks 0-6

- 1) Hand, wrist and elbow active motion
- 2) Active scapular motion (shrugs, retraction)

### Phase 2

Weeks 7-9

- 1) Begin active shoulder motion in all planes – (no restrictions unless otherwise specified)
  - Forward elevation
  - External rotation
  - Extension
  - Abduction
- 2) Initiate independent home motion  
Home exercise program - AAROM for external rotation, wall climbs

### Phase 3

Weeks 10-12

- 1) Continue motion as needed
- 2) Begin Cuff Strengthening when approved
- 3) Progress to overhead resistance with proper scapulohumeral rhythm
- 3) Avoid external rotation, 90/90, or throwing position

Week 13+

- 1) Advance activity
  - Swimming (avoid extreme abduction/external rotation)
  - Begin throwing program progression if approved
  - No contact or at risk activities/sports until 6 months

6 months

- 1) Return to regular activity
- 2) Maintenance Program
  - Daily Stretching (all planes)
  - Strengthening 2-3 times / week

